

1 Arden's rule

Exercice 1. Let \mathcal{A}, \mathcal{B} be regular langages, prove that, if $\epsilon \notin \mathcal{A}$ then $\mathcal{X} = \mathcal{A}\mathcal{X} \cup \mathcal{B}$ has a unique language \mathcal{X} solution. Express this solution in terms of a (resp. b) a regular expression recognizing \mathcal{A} (resp. \mathcal{B}).

Advice :

1. find a first solution ;
2. show that any solution is larger than your solution ;
3. show that any solution is contained within your solution.

2 Pumping lemma

Exercice 2. Justify whether or not the following languages are regular ?

1. $\{a^n b^n \mid n \in \mathbb{N}\}$
2. $\{a^m b^n \mid n \equiv m \pmod{d}\}$ for a given $d \in \mathbb{N}$.
3. $\{a^p \mid p \text{ premier}\}$
4. $\{a^{P(n)} \mid n \in \mathbb{N}\}$ for a given $P \in \mathbb{N}[X]$.

3 Puzzles

Exercice 3. Let x and y be two words such that $xy = yx$. What can you say about the shape of x and y ?

Exercice 4. Let $(p, q) \in \mathbb{N}^*$, and \mathcal{L} a regular language. When is $\frac{p}{q}\mathcal{L} = \left\{u \mid \exists v : uv \in \mathcal{L} \text{ et } |u| = \frac{p}{q}|uv|\right\}$ also a regular language ?

4 Generalized Arden's rule

Definition 1. A monoid is an algebraic structure (E, \odot, e) such that :

- (binary operation) For $x, y \in E^2$, $x \odot y \in E$
- (associative) For $x, y, z \in E^3$, $(x \odot y) \odot z = x \odot (y \odot z)$
- (identity element) For $x \in E$, $x \odot e = e \odot x = x$

Definition 2. A semiring is an algebraic structure $(E, +, \times, 0, 1)$ such that :

- $(E, +, 0)$ is a commutative monoid ($a + b = b + a$);
- $(E, \times, 1)$ is a monoid;
- \times is distributive over $+$ ($a \times (b + c) = a \times b + a \times c$ and $(b + c) \times a = b \times a + c \times a$);
- 0 is absorbing for \times (i.e. $0 \times a = a \times 0 = 0$)

Exercice 5. Notice that $(\mathcal{L}_\Sigma, \cup, /, \emptyset, \epsilon)$ is a semiring where :

- \mathcal{L}_Σ is the set of regular languages over Σ ;
- $/$ is the concatenation $X/Y = \{xy \mid x \in X, y \in Y\}$;
- \cup is the union;
- \emptyset is the empty language;
- ϵ is the language containing only the empty word.

Let $(M_n(\mathcal{L}_\Sigma), +, \times, 0, 1)$ be the set of square matrices of size n over the semiring $(\mathcal{L}_\Sigma, /, \cup, \emptyset, \epsilon)$ with

- $(M + N)_{i,j} = M_{i,j} \cup N_{i,j}$;
- $(M \times N)_{i,j} = \bigcup_{1 \leq k \leq n} M_{i,k} / N_{k,j}$;
- $0_{i,j} = \emptyset$

$$- 1_{i,j} = \begin{cases} \epsilon & i = j \\ \emptyset & \text{sinon} \end{cases}$$

$$- M^* = \sum_{i \in \mathbb{N}} M^k$$

Notice that $(M_n(\mathcal{L}_\Sigma), +, \times, 0, 1)$ is also a semiring.


Remark : any semiring can be transformed into a semiring of matrices

Exercice 6. Let $\mathcal{A} \in M_n(\mathcal{L}_\Sigma)$ with $\forall i, j \in \mathcal{A}_{i,j}$ and $\mathcal{B} \in M_{n,1}(\mathcal{L}_\Sigma)$ a vector of size n , show that there is a unique solution to $\mathcal{X} \in M_{n,1}(\mathcal{L}_\Sigma)$ with $\mathcal{X} = \mathcal{A}\mathcal{X}^T \cup \mathcal{B}$ (i.e. $\forall i : \mathcal{X}_i = (\bigcup_j \mathcal{A}_{i,j}/\mathcal{X}_j) \cup \mathcal{B}_i$).

Exercice 7. (*Gaussian elimination*) Given $\mathcal{A} \in M_n(\mathcal{L}_\Sigma)$ and $\mathcal{B} \in M_{n,1}(\mathcal{L}_\Sigma)$ show that when $n > 1$ the languages $(\mathcal{A}^* \mathcal{B})_i$ for $i \in \{1, \dots, n - 1\}$ can be expressed as $\mathcal{A}'^* \mathcal{B}'_i$ for a given $\mathcal{A}' \in M_n(\mathcal{L}_\Sigma), \mathcal{B}' \in M_{n,1}(\mathcal{L}_\Sigma)$.

Exercice 8. Deduce an algorithm returning the regular expression corresponding to a given automata.

Exercice 9. What is the regular expression corresponding to :



5 Monoid of a language

Definition 3. Given an alphabet Σ and a language \mathcal{L} (not necessarily regular) we state $\mu_{\mathcal{L}}$ le syntactic morphism from \mathcal{L} defined as : $\mu_{\mathcal{L}}(u) = \{(x, y) \in \Sigma^{*2} \mid xuy \in \mathcal{L}\}$.

Exercice 10. Show that $\mu_{\mathcal{L}}$ is compatible with the concatenation, i.e. $\forall u, v, w : \mu_{\mathcal{L}}(v) = \mu_{\mathcal{L}}(w) \Rightarrow (\mu_{\mathcal{L}}(vu) = \mu_{\mathcal{L}}(wu)) \wedge (\mu_{\mathcal{L}}(uv) = \mu_{\mathcal{L}}(uw))$.

Exercice 11. For each element $e \in \mu_{\mathcal{L}}(\Sigma^*)$ we chose a an element $r(e)$ tel que $\mu_{\mathcal{L}}(r(e)) = e$, and we define $u \odot v = \mu_{\mathcal{L}}(r(u)) \odot \mu_{\mathcal{L}}(r(v)) = \mu_{\mathcal{L}}(r(u)r(v))$.

Justify that :

- \odot does not depend on the choices for r .
- the structure $(\mu_{\mathcal{L}}(\Sigma^*), \odot, \mu_{\mathcal{L}}(\epsilon))$ induced by $\mu_{\mathcal{L}}$ is a monoid

and thus $\mu_{\mathcal{L}}$ do is a morphism from the the free monoid over Σ towards the monoid induced by $\mu_{\mathcal{L}}$.

Remark : any monoid (M, \odot, e) compatible with a function μ induce a structure of monoid over $\mu(M)$.

Exercice 12. Prove that $\mu_{\mathcal{L}}(\Sigma^*)$ is finite when \mathcal{L} is regular.

Exercice 13. Prove that when $\mu_{\mathcal{L}}(\Sigma^*)$ is finite then \mathcal{L} is regular.

Exercice 14. What does the the syntactic monoid of the well-parenthesized words looks like ?